

Title	Neogene climate of NW Africa		
Proponents	Torsten Bickert, Ilham Bouimetarhan, Anya J. Crocker, Peter deMenocal, Timothy D. Herbert, Sebastian Krastel, Anna Nele Meckler, Stefan Mulitza, Oscar Romero, Enno Schefuß, Tilmann Schwenk, Thomas Westerhold, Paul A. Wilson		
Keywords	NW-African-continental-margin, Neogene-hydroclimate, Sahara, eastern-boundary upwelling	Area	NW African continental margin

Proponent Information

Proponent	Torsten Bickert
Affiliation	MARUM, University of Bremen, 28359 Bremen, Germany
Country	Germany

Permission is granted to post the coversheet/site table on www.iodp.org

Abstract

We propose an IODP expedition to investigate the Miocene to Quaternary climate along a latitudinal transect of drill sites from the continental margin of NW Africa from Cap Bojador (~28°N) southeast of the Canary Islands to Sierra Leone Rise (~5°N). Major objectives at all primary sites are (i) to study North African hydroclimate and vegetation response to global and North Atlantic warmth, and (ii) to reconstruct the timing and cause of Neogene aridification and linkages to global climate state and ocean circulation. These new archives will complement those recently recovered from Southern Africa by IODP Exp. 361, and from International Continental Scientific Drilling Program coring of East African lakes, to build an in-depth understanding of past natural climate variability under future-relevant conditions across Africa. A related secondary objective is to study ocean productivity and ecosystem coupling to African hydroclimate and global and North Atlantic warmth. Our drilling strategy is based on an extensive array of high quality multichannel seismic data combined with information from GeoB sediment cores and the results of DSDP/ODP Legs 41 and 108. Drill cores from those two early legs provide helpful stratigraphic control and highlight the potential of our proposed latitudinal transect to transform our understanding of rainfall response in North Africa to changes in global climate state. Our plans include two sites positioned close to classic sites (366 on Sierra Leone Rise and 659 on Cape Verde Plateau) where drilling will recover the more complete sections needed, and a further three primary and nine alternate sites where new coring will provide unprecedented high-quality information on African rainfall and climate variability and vegetation response as well as oceanic production along the North African margin. Our scientific objectives are closely aligned with those of the IODP 2013-2023 Science Plan, and the proposed expedition will help meet the information priorities for human adaption to climate change in NW Africa identified by IPCC 2013 working group 2.

Scientific Objectives

Objective 1 (primary priority) - NW African climate in a warmer world, with emphasis on wind systems, dust supply, precipitation and vegetation

- How did African climate and vegetation respond to global warmth during the late Cenozoic? What are the lessons from these records of natural climate variability for the Anthropocene?
- What is the relationship between NW African hydroclimate, global warmth and Atlantic overturning ocean circulation?
- How old is the world's largest hot desert, the Sahara, and what forcing mechanisms controlled its origin and evolution?

Objective 2 (secondary priority) - Ocean productivity and ecosystem response to climate perturbations

- What is the marine productivity response to climate conditions warmer than today?
- What is the role of dust in fertilizing marine bioproduction and ballasting particle fluxes?
- What are the consequences of future climate change for marine ecosystems off NW Africa?

Non-standard measurements technology needed to achieve the proposed scientific objectives

Proposed Sites (Total proposed sites: 14; pri: 5; alt: 9; N/S: 0)

Site Name	Position (Lat, Lon)	Water Depth (m)	Penetration (m)			Brief Site-specific Objectives
			Sed	Bsm	Total	
NWAFR-01A (Alternate)	27.5374 -13.8459	1430	500	0	500	Cape Bojador high-resolution climate, Plio/Pleistocene millennial scale African hydroclimate, 0-3,5 Ma
NWAFR-02B (Primary)	24.2663 -17.1061	1590	1100	0	1100	Early Miocene climate, Sahara summer dust input, Neogene northern limit of Mauritania upwelling, 0 to 17-20 Ma
NWAFR-03B (Alternate)	24.2694 -17.1585	1690	1060	0	1060	Early Miocene climate, Sahara summer dust input, Neogene northern limit of Mauritania upwelling, 0 to 17-20 Ma
NWAFR-04A (Alternate)	20.85154 -18.41738	2140	500	0	500	Cape Blanc high-resolution climate and productivity, Plio/Pleistocene millennial scale African hydroclimate, 0 to 3.5 Ma
NWAFR-05B (Primary)	15.4865 -17.9131	2280	250	0	250	North Senegal high-resolution climate, Pleistocene millennial scale African hydroclimate, 0-1,5 Ma
NWAFR-06B (Primary)	13.5269 -18.4441	3540	500	0	500	Early Miocene African climate, South Senegal high-resolution hydroclimate, 0-16 Ma
NWAFR-07A (Alternate)	18.0772 -21.0262	3069	300	0	300	Sahara dust summer plume, early Miocene African climate, 0-20 Ma
NWAFR-08B (Primary)	5.7587 -20.0084	2880	250	0	250	Sahara dust winter plume, Early Miocene African climate, 0-19 Ma
NWAFR-09B (Alternate)	27.5376 -13.7022	900	650	0	650	Cape Bojador high-resolution climate, Plio/Pleistocene millennial scale African hydroclimate, 0-3,0 Ma
NWAFR-10B (Alternate)	20.5454 -18.0512	900	550	0	550	Cape Blanc high-resolution climate and productivity, Plio/Pleistocene millennial scale African hydroclimate, 0 to 4.5 Ma
NWAFR-11B (Alternate)	12.4293 -18.0054	3001	300	0	300	Casamanche high-resolution climate, Plio/Pleistocene millennial scale African hydroclimate, 0-4,0 Ma
NWAFR-12C (Alternate)	18.7383 -21.0029	3180	300	0	300	Sahara dust summer plume, early Miocene African climate, 0-20 Ma
NWAFR-13A (Alternate)	5.6742 -19.8437	2865	250	0	250	Sahara dust winter plume, Early Miocene African climate, 0-19 Ma
NWAFR-14A (Primary)	17.9509 -20.8328	2850	300	0	300	Sahara dust summer plume, early Miocene African climate, 0-20 Ma